


IN THE CLAIMS:

- Sub C2 / 1. (AMENDED). A stabilized pharmaceutical preparation, comprising: (a) an α amino acid; (b) an optional auxiliary agent for manufacturing a pharmaceutical preparation; and (c) a 4-amino-3-substituted-butanoic acid derivative, which 4-amino-3-substituted-butanoic acid derivative has the general formula:



wherein,

R₁ is a hydrogen atom, a hydroxyl group, a methyl group or an ethyl group;

R₂ is a monovalent group selected from:

a straight or branched alkyl group of 3 - 8 carbon atoms;

a straight or branched alkylene group of 3 - 8 carbon atoms;

A2 a straight or branched alkyl group of 3 - 8 carbon atoms which is mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 3 - 8 carbon atoms;

a cycloalkyl group of 3 - 8 carbon atoms which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group,

h
C
cont

an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedieryl group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedieryl group of 5 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms, is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, and one or two of the unsubstituted methylene groups (-CH₂-) are mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

Ag

Sub
C2
cont

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, and one or two of the unsubstituted methylene groups (-CH₂-) being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

Az
a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, said phenyl group being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or

Sub
C2
cont

-S(O)₂-, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, one of the methylene groups (-CH₂-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)₂;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, and one of the methylene groups (-CH₂-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)₂- and one or two of the unsubstituted methylene groups (-CH₂-) being mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a phenyl or naphthyl group;

a phenyl group substituted with a methylenedioxy group;

a phenyl or naphthyl group which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group, a phenoxy group, a phenylmethoxy group, a phenylmethoxy group wherein said phenyl ring is mono-substituted with a halogen atom, trifluoromethyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring, a cycloalkanedienylmethoxy group having 5 - 8 carbon atoms in the cycloalkanedienyl ring, a cycloalkylmethoxy group wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring having 5 - 8 carbon atoms is

A2

Sub
C2
cont

replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, a cycloalkenylmethoxy group wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, a cycloalkanedieryl-methoxy group wherein one of the methylene groups (-CH₂-) in said cycloalkanedieryl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂- group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring wherein said cycloalkyl ring is mono-substituted with a halogen atom, trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring wherein said cycloalkenyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH₂-) in said cycloalkenyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, or a cycloalkanedierylmethoxy group having 5 - 8 carbon atoms in the cycloalkanedieryl ring wherein said cycloalkanedieryl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH₂-) in said cycloalkanedieryl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS-;

an -O-, -S- or -SS-phenyl group;

Az

Sub
C2
cont

a diphenylamino group;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an -O-, -S- or -SS-phenyl group wherein said phenyl group is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

or

R₁ and R₂, together with the carbon atom to which they are attached, may form a divalent group selected from:

a cycloalkylidene group of 5 - 8 carbon atoms;

a cycloalkylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group or a carboxyl group;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂- group and one or more of the unsubstituted methylene groups

Az

Sub
C2
cont

(-CH₂-) in said cycloalkyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedierylidene group of 5 - 8 carbon atoms;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedierylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

A₂
a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedierylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedieryl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedierylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedieryl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂- group and one or more of the unsubstituted methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedieryl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms, said phenyl ring being mono-, di-,

Sub
C2
cont

tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group; or

provided that when R₂ is a phenyl or naphthyl group which is mono-, di- or tri-substituted with a halogen atom, the α -amino acid is not glycine.

2. (AMENDED). The stabilized pharmaceutical preparation of Claim 1 wherein said α -amino acid is one or more selected from

the L-, D- and DL-forms of neutral α -amino acids;

alkali salts, acid amides, alkyl-substituted derivatives of acid amides or alkyl esters of the L-, D- and DL-forms of acidic α -amino acids;

acid addition salts or monoacylated derivatives of the L-, D- and DL-forms of basic α -amino acids;

α,ω -diaminodicarboxylic acids; and

acidic amino acid-basic amino acid adducts of the L-, D- and DL-forms of acidic α -amino acids and the L-, D- and DL-forms of basic α -amino acids.

3. (AMENDED). The stabilized pharmaceutical preparation of Claim 2 wherein said α -amino acid is one or more selected from

Sub
C1

neutral α -amino acids consisting of glycine, phenylglycine, hydroxyphenylglycine, dihydroxyphenylglycine, L-alanine, hydroxy-L-alanine, L-leucine, hydroxy-L-leucine, dihydroxy-L-leucine, L-norleucine, methylene-L-norleucine, L-ketonorleucine, L-isoleucine, hydroxy-L-isoleucine, dihydroxy-L-isoleucine, L-valine, hydroxy-L-valine, L-isovaline, L-norvaline, hydroxy-L-norvaline, hydroxy-L-ketonorvaline, L-methionine, L-homomethionine, L-ethionine, L-threonine, acetyl-L-threonine, L-tryptophan, hydroxy-L-tryptophan, methyl-L-tryptophan, L-tyrosine, hydroxy-L-tyrosine, methyl-L-tyrosine, bromo-L-tyrosine, dibromo-L-tyrosine, 3,5-diiodo-L-tyrosine, acetyl-L-tyrosine, chloro-L-tyrosine, L-m-tyrosine, L-levodopa, L-methyldopa, L-thyroxine, L-serine, acetyl-L-serine, L-homoserine, acetyl-L-homoserine, ethyl-L-homoserine, propyl-L-homoserine, butyl-L-homoserine, L-cystine, L-homocystine, methyl-L-cysteine, allyl-L-cysteine, propyl-L-cysteine, L-phenylalanine, dihydro-L-phenylalanine, hydroxymethyl-L-phenylalanine, L-aminobutyric acid, L-aminoisobutyric acid, L-ketoaminobutyric acid, dichloro-L-aminobutyric acid, dihydroxy-L-aminobutyric acid, phenyl-L-aminobutyric acid, L-aminovaleric acid, L-aminohydroxyvaleric acid, dihydroxy-L-aminovaleric acid, L-aminoisovaleric acid, L-aminohexanoic acid, methyl-L-aminohexanoic acid, L-aminoheptanoic acid, L-aminooctanoic acid and citrulline and the D- and DL-forms thereof;

acidic α -amino acids consisting of L-aspartic acid, L-glutamic acid, L-carbocysteine, L-aminoglutaric acid, L-aminosuccinic acid, L-aminoadipic acid, L-aminopimelic acid, hydroxy-L-aminopimelic acid, methyl-L-aspartic acid, hydroxy-L-aspartic acid, methyl-L-glutamic acid, methyl-hydroxy-L-glutamic acid, L-methyleneglutamic acid, hydroxy-L-glutamic acid, dihydroxy-L-glutamic acid and hydroxy-L-aminoadipic acid and the D- and DL-forms thereof;

basic α -amino acids consisting of L-arginine, L-lysine, L-ornithine, L-canavanine, L-canaline, hydroxy-L-lysine, L-homoarginine, hydroxy-L-homoarginine, hydroxy-L-ornithine, L-diaminopropionic acid, L-diaminohexanoic acid, L-diaminobutyric acid, L-diaminovaleric acid, L-diaminoheptanoic acid, and L-diaminooctanoic acid and the D- and DL-forms thereof; and

α,ω -diaminodicarboxylic acids consisting of diaminosuccinic acid, diaminoglutaric acid, diaminoadipic acid and diaminopimelic acid;

provided that, when said α -amino acid is an aliphatic α -amino acid, it is used in the form of the corresponding alkali salt, acid amide, alkyl-substituted derivative of acid amide or alkyl ester thereof, or

when said α -amino acid is a basic α -amino acid, it is used in the form of the corresponding acid addition salt or monoacylated derivative thereof, or

said acidic α -amino acid and said basic α -amino acid are also used in the form of the corresponding acidic amino acid-basic amino acid adduct.

- Sub
B1
A2
4. (AMENDED). A stabilized pharmaceutical preparation as claimed in Claims 1, 2, or 3 wherein a total amount of said α -amino acid is in the range of 0.001 - 80 moles per mole of the 4-amino-3-substituted-butanoic acid derivative.
 5. (AMENDED). A stabilized pharmaceutical preparation as claimed in Claims 1, 2, 3, or 4 wherein it is in the form of liquid preparations.
 6. (AMENDED). A stabilized pharmaceutical preparation as claimed in Claim 5 wherein it is in the dosage form of liquid preparations, syrups or injections.
 7. (AMENDED). A stabilized pharmaceutical preparation as claimed in Claims 1, 2, 3, or 4 wherein it is in the form of solid preparations.
 8. (AMENDED). A stabilized pharmaceutical preparation as claimed in Claim 7 which is in the dosage form of tablets, powders, granules or capsules.
 9. (AMENDED). A stabilized pharmaceutical preparation as claimed in Claims 1, 2, 3, 4, 5, 6, 7, or 8 which is a gabapentin-containing preparation, a pregabalin-containing preparation, a baclofen-containing preparation, or a preparation containing 3-aminomethyl-4-cyclohexyl-butanoic acid, 3-aminomethyl-5-cyclohexyl-pentanoic

A₂ sub 11
acid, 3-aminomethyl-4-phenyl-butanoic acid or 3-aminomethyl-5-phenyl-pentanoic acid, provided that the baclofen-containing preparation does not contain glycine as the α -amino acid.

- A₃ sub 12
18. (NEW). A stabilized pharmaceutical preparation containing an α amino acid, an auxiliary agent for manufacturing a pharmaceutical preparation if necessary, and gabapentin.
19. (NEW). The stabilized pharmaceutical preparation of Claim 18 wherein the α amino acid is as provided in Claims 2 or 3.
20. (NEW). The stabilized pharmaceutical preparation containing of Claims 18 or 19 in the form of liquid preparations, syrups, or injections.
21. (NEW). The stabilized pharmaceutical preparation of Claims 18 or 19 in the form of solid preparations.
22. (NEW). The stabilized pharmaceutical preparation of Claim 21 in the dosage forms of tablets, powders, granules, or capsules.